## Exercise 9: Stabilization

## Task 1: Recovering Reasonable Constraints

In this exercise, we want to get a better handle on the set of constraints from the lecture:

$$S(1) \ge 2\left(\delta + \left(1 - \frac{1}{\vartheta}\right)T\right) \tag{1}$$

$$\frac{R^{-}}{\vartheta} \ge \sigma_h + \vartheta \mathcal{S}(1) + d \tag{2}$$

$$\frac{B_2}{\vartheta} > \sigma_h + R^+ + T + 2\mathcal{S}(1) \tag{3}$$

$$B_1 > \sigma_h + R^+ \tag{4}$$

$$B_3 > R^+ + (M-1)(T+S(1)) + (\vartheta+1)S(M) + \sigma_h$$
 (5)

$$B_2 \le \frac{R^-}{\vartheta} + (M-1)\left(\frac{T}{\vartheta} - \mathcal{S}(1)\right) + \mathcal{S}(M) \tag{6}$$

$$\frac{R^+}{\vartheta} \ge (\vartheta + 1)\mathcal{S}(M) + \sigma_h \tag{7}$$

$$2(\mathcal{S}(1) - \mathcal{S}(M)) \ge \sigma_h \tag{8}$$

- a) Choose  $R^-$  tight according to (2) and  $R^+/\vartheta$  equal to the r.h.s. of (7) plus d, respectively. Show that for these choices (1) and (8) hold. (Hint: Use these equalities to express  $\mathcal{S}(1)$  in terms of  $\mathcal{S}(M)$  and other terms not involving  $R^+$  or  $R^-$ .)
- b) Fix these choices and consider the remaining inequalities. Which of the terms on the r.h.s. of the inequalities are in  $\mathcal{O}(\sigma_h + d)$ , if M is not treated as a constant? (Hint: Recall that  $\vartheta \in \mathcal{O}(1)$ .)
- c) In addition, suppose now that for  $\alpha > 1$ , we have that  $B_3 = \alpha B_2 = \alpha^2 B_1$  and can choose  $B_1 > 0$  sufficiently large. For which values of  $\alpha$  can you choose M so that the system of inequalities is satisfied?

## Task 2: If it Were so Simple...

- a) Modify the Srikanth-Toug algorithm so that its pulses can be triggered by an external NEXT signal in the vein of Definition 9.7. (Hint: Add a second, smaller timeout to the transition from READY to PROPOSE.)
- b) What choices of  $B_1$ ,  $B_2$ , and  $B_3$  can this solution support?