



Karl Bringmann and Marvin Künnemann

Winter 2017/18

Exercises for Fine-Grained Complexity Theory

www.mpi-inf.mpg.de/departments/algorithms-complexity/teaching/winter17/fine-complexity/

Exercise sheet 1

Due: **Tuesday, November 7, 2017**

Total points : 40

You are allowed to collaborate on the exercise sheets, but you have to write down a solution on your own, **using your own words**. Please indicate the names of your collaborators for each exercise you solve. Further, cite all external sources that you use (books, websites, research papers, etc.).

You need to collect at least 50% of all points on exercise sheets.

Exercise 1 (10 points) Let $0 < \varepsilon < 0.5$ and $c \in \mathbb{N}$. Consider the following functions:

$$n^{2-\varepsilon}, \quad n \cdot \log^c n, \quad n, \quad \frac{n^2 \log \log n}{\log n}, \quad \frac{n^2}{2^{\sqrt{\log n}}}, \quad n^2, \quad n^{1+\varepsilon}, \quad n^{2-\frac{1}{\log \log n}}.$$

- (8 points) Order these functions according to their asymptotic growth. Justify your answers.
- (2 points) Which of these functions are contained in $n^{1+o(1)}$ and $n^{2-o(1)}$, respectively?

Exercise 2 (10 points) In the lecture we introduced the *Orthogonal Vectors Hypothesis*:

OVH: Given two sets $A, B \subseteq \{0, 1\}^d$ such that $|A| = |B| = n$. There is no algorithm running in time $O(n^{2-\varepsilon} \cdot \text{poly}(d))$ (for any $\varepsilon > 0$) which decides whether there exists $a \in A, b \in B$ such that a and b are orthogonal.

- (5 points) Consider the following variant **UOVH** of **OVH**:

UOVH (Unbalanced **OVH**): Given two sets $A, B \subseteq \{0, 1\}^d$ such that $|A| = n, |B| = \sqrt{n}$. There is no algorithm running in time $O(n^{1.5-\varepsilon} \cdot \text{poly}(d))$ (for any $\varepsilon > 0$) which decides whether there exist $a \in A, b \in B$ such that a and b are orthogonal.

Prove that **UOVH** and **OVH** are equivalent.

- (5 points) Consider the following stronger variant **LDOVH** of **OVH**:

LDOVH (Low-dimensional **OVH**): For all $\varepsilon > 0$ there is a (constant) $c > 0$ such that the Orthogonal Vectors problem cannot be solved in time $O(n^{2-\varepsilon})$ even restricted to $d \leq c \cdot \log n$.

Prove that **SETH** implies **LDOVH**.

(Hint: You may use the Sparsification Lemma from the lecture.)

Exercise 3 (10 points) One of the classical string problems is finding the longest common substring of two given strings A and B . In this exercise we will consider a variation of that problem, where we allow one string to contain *wildcard* characters, i.e., characters that are considered equal to any other character. Formally, consider the following problem:

Longest Common Substring With Don't Cares: Given a string A of length n over some alphabet Σ and string B of length n over the alphabet $\Sigma \cup \{\star\}$, find the length of the longest string that is a substring of both A and B , where a “ \star ” in B can be treated as any character from the alphabet Σ .

Prove that there is no $O(n^{2-\varepsilon})$ time algorithm (for any $\varepsilon > 0$) for this problem unless **OVH** fails.

Exercise 4 (10 points) Prove that **SETH** implies **ETH**.

(Hint: You may use the Sparsification Lemma from the lecture.)